

Deformations of holomorphic Lagrangian fibrations*

Justin Sawon

August, 2005

Abstract

Let $X \rightarrow \mathbb{P}^n$ be a $2n$ -dimensional projective holomorphic symplectic manifold admitting a Lagrangian fibration over $\mathbb{P}^n$. Matsushita proved that the fibration can be deformed in a codimension one family in the moduli space $\text{Def}(X)$ of deformations of X . We extend his result by proving that if the Lagrangian fibration admits a section, then there is a codimension two family of deformations which also preserve the section.

1 Introduction

Let X be a $2n$ -dimensional compact Kähler manifold. We say X is a *holomorphic symplectic manifold* if it admits a closed two-form σ of type $(2, 0)$ which is non-degenerate in the sense that $\sigma^{\wedge n}$ trivializes the canonical bundle $K_X = \Omega^{2n}$. Moreover, we call X *irreducible* if $H^0(X, \Omega^2)$ is one-dimensional and generated by $[\sigma]$. Huybrechts' notes in [7] provide a comprehensive introduction to the standard results on irreducible holomorphic symplectic manifolds.

Let us summarize what is known about fibrations on X . Suppose we have a proper (holomorphic) surjection $f : X \rightarrow B$ onto a complex space B such that the general fibre is connected and $0 < \dim B < 2n$. If X is projective and B is a normal variety then Matsushita [9, 10] showed that

- every irreducible component of a fibre of f is a (holomorphic) Lagrangian submanifold of X ; in particular, it is n -dimensional,
- the generic fibre is an abelian variety,
- B is n -dimensional and has only $\mathbb{Q}$ -factorial log-terminal singularities,
- K_B^* is ample and B has Picard number one.

*2000 *Mathematics Subject Classification.* 53C26; 14D06; 14J60.

Moreover, if B is smooth then it has the same Hodge numbers as $\mathbb{P}^n$ (see [11]). In particular, for $n = 2$ the base is isomorphic to $\mathbb{P}^2$ (a similar result was obtained by Markushevich [8]). In general it is expected that B should be isomorphic to $\mathbb{P}^n$, and more evidence is provided by a result of Cho, Miyaoka, and Shepherd-Barron [3] which states that if $f : X \rightarrow B$ admits a section (or even local sections) then $B \cong \mathbb{P}^n$. Huybrechts [7, Proposition 24.8] extended some of these results by dropping the projectivity assumption: he showed that if X and B are (smooth) Kähler manifolds then

- every fibre of f is (holomorphic) Lagrangian,
- every smooth fibre is a complex torus,
- B is n -dimensional and projective,
- K_B^* is ample and its Picard and second Betti numbers both equal one.

In this paper we consider deformations of $f : X \rightarrow B$. It is known that the Kuranishi space $\text{Def}(X)$ of deformations of X is a smooth complex manifold of dimension $b_2 - 2$, where b_2 is the second Betti number of X . Under the assumptions that X is projective and $B \cong \mathbb{P}^n$, Matsushita [11] proved that there is a codimension one submanifold of $\text{Def}(X)$ parametrizing deformations of X which are Lagrangian fibrations over $\mathbb{P}^n$. We include a proof of this result in Section 3 in order to set-up our notation. Our main result is Theorem 6: we prove that if X is projective and $f : X \rightarrow B$ admits a section, then there is a codimension two submanifold of $\text{Def}(X)$ parametrizing deformations which are Lagrangian fibrations and admit sections.

Let X' be an arbitrary irreducible holomorphic symplectic manifold. Can X' be deformed to a Lagrangian fibration? The answer in general is unknown. However, the question has been answered in the affirmative for all known examples of irreducible holomorphic symplectic manifolds: see Beauville [2] for the Hilbert schemes of points on a K3 surface, Debarre [4] for the generalized Kummer varieties, and Rapagnetta [12] for O'Grady's examples.

The author would like to thank Akira Fujiki, Daniel Huybrechts, and Daisuke Matsushita for useful conversations. Thanks also to Ryushi Goto for the invitation to Japan where this result was obtained. The author is supported by NSF grant number 0305865.

2 Some preliminaries

We collect together some results that we will need later.

Theorem 1 (Matsushita [11]) *Let $f : X \rightarrow B$ be a Lagrangian fibration between smooth projective manifolds. Then*

$$R^j f_* \mathcal{O}_X \cong \Omega_B^j.$$

Let X be a projective irreducible holomorphic symplectic manifold which admits a Lagrangian fibration $f : X \rightarrow \mathbb{P}^n$ over projective space. Let $L := f^*\mathcal{O}(1)$ be the pull-back of the hyperplane line bundle to X . We can use Theorem 1 to calculate the cohomology of L .

Lemma 2 *For (X, L) as above*

$$h^0(X, L) := \dim H^0(X, L) = n + 1$$

and all higher cohomology of L vanishes.

Proof The Leray spectral sequence gives

$$H^i(\mathbb{P}^n, R^j f_* L) \Rightarrow H^{i+j}(X, L).$$

On the left hand side

$$\begin{aligned} R^j f_* L &= R^j f_*(f^*\mathcal{O}(1)) \\ &\cong \mathcal{O}(1) \otimes R^j f_* \mathcal{O}_X \\ &\cong \mathcal{O}(1) \otimes \Omega_{\mathbb{P}^n}^j \end{aligned}$$

by the projection formula and Theorem 1. We can therefore use Bott's generalization of the Borel-Weil theorem [5] to calculate the spectral sequence, finding

$$H^i(\mathbb{P}^n, \mathcal{O}(1) \otimes \Omega_{\mathbb{P}^n}^j) \cong \mathbb{C}^{n+1}$$

when $i = j = 0$, and vanishes otherwise. Therefore the sequence collapses, and the lemma follows. $\square$

Remark The proof of the lemma relies on two important assumptions: that the base of the fibration is $\mathbb{P}^n$ and that X is projective (note that the projectivity of X is crucial for Theorem 1).

We need to know how the cohomology of a line bundle behaves in a family.

Theorem 3 (Grauert and Remmert [6], page 210) *Let $\mathcal{X} \rightarrow \Delta$ be a deformation of a complex manifold $X = \mathcal{X}_0$, where Δ is a polydisc of some dimension around 0. Let V be a vector bundle on $\mathcal{X}$, with restriction V_t to each $\mathcal{X}_t$. Then for $i \geq 0$,*

$$h^i(\mathcal{X}_t, V_t) := \dim H^i(\mathcal{X}_t, V_t)$$

is an upper semi-continuous function of $t \in \Delta$. In other words

$$A_{i,d} := \{t \in \Delta \mid h^i(\mathcal{X}_t, V_t) \geq d\}$$

is a closed analytic subset of Δ for each integer d .

3 Deforming fibrations

Let X be an irreducible holomorphic symplectic manifold of dimension $2n$. For the following statements, see Section 22 of Huybrechts' notes in [7]. Denote by

$$\mathcal{X} \rightarrow (\text{Def}(X), 0)$$

the Kuranishi family parametrizing local deformations of $X = \mathcal{X}_0$. We think of $(\text{Def}(X), 0)$ as the germ of a complex space. It is smooth (deformations are unobstructed) of dimension $b_2 - 2$, where b_2 is the second Betti number of X . Note that when we deform X as a complex manifold it remains holomorphic symplectic and irreducible; for small deformations it also stays Kähler.

The germ $(\text{Def}(X), 0)$ can be represented by a contractible open set, and therefore for each $t \in \text{Def}(X)$ we can choose an isomorphism

$$\psi_t : H^2(\mathcal{X}_t, \mathbb{Z}) \rightarrow H^2(X, \mathbb{Z})$$

known as a *marking*. Let

$$Q_X := \{[\alpha] \in \mathbb{P}(H^2(X, \mathbb{C})) \mid q_X(\alpha) = 0 \text{ and } q_X(\alpha + \bar{\alpha}) > 0\}$$

be the *period domain* of X , where q_X is the Beauville-Bogomolov quadratic form on $H^2(X, \mathbb{Z}) \otimes \mathbb{C}$. Then by the Local Torelli Theorem (see Beauville [1]), the *period map*

$$\mathcal{P}_X : (\text{Def}(X), 0) \rightarrow (Q_X, [\sigma])$$

which takes t to $[(\psi_t \otimes \mathbb{C})(\sigma_t)]$ is a local isomorphism.

In this section we prove the following result.

Theorem 4 (Matsushita [11], Corollary 1.7) *Let X be an irreducible projective holomorphic symplectic manifold which admits a Lagrangian fibration $f : X \rightarrow \mathbb{P}^n$ over projective space. There is a codimension one submanifold $\Delta^f \subset \text{Def}(X)$ (containing zero) which parametrizes holomorphic symplectic manifolds X' which admit Lagrangian fibrations over $\mathbb{P}^n$.*

Proof Let $L := f^*\mathcal{O}(1)$ be the pull-back of the hyperplane line bundle to X . Then f is clearly given by the morphism

$$\phi_L : X \rightarrow \mathbb{P}(H^0(X, L)^*)$$

induced by the linear system of L . Our aim is to extend L and this map to certain deformations of X .

Let $c_1 := c_1(L) \in H^2(X, \mathbb{Z}) \cap H^{1,1}(X)$ be the first Chern class of L , and let

$$\Delta^f := \{t \in \text{Def}(X) \mid \psi_t^{-1}(c_1) \in H^2(\mathcal{X}_t, \mathbb{Z}) \text{ is of type } (1, 1)\}.$$

If $t \in \Delta^f$, then $\psi_t^{-1}(c_1)$ is orthogonal to σ_t with respect to $q_{\mathcal{X}_t}$, or equivalently, c_1 is orthogonal to $(\psi_t \otimes \mathbb{C})(\sigma_t)$ with respect to q_X . Therefore Δ^f maps isomorphically to a neighbourhood of $[\sigma]$ in

$$Q_X^f := \{[\alpha] \in Q_X \mid q_X(\alpha, c_1) = 0\}$$

which is of codimension one in Q_X . Hence Δ^f must be codimension one in $\text{Def}(X)$.

The exponential exact sequence on $\mathcal{X}_t$ gives

$$\dots \rightarrow 0 \rightarrow H^1(\mathcal{X}_t, \mathcal{O}^*) \rightarrow H^2(\mathcal{X}_t, \mathbb{Z}) \xrightarrow{\beta} H^2(\mathcal{X}_t, \mathcal{O}) \rightarrow \dots$$

since $H^1(\mathcal{X}_t, \mathcal{O})$ vanishes. For $t \in \Delta^f$, $\psi_t^{-1}(c_1)$ is of type $(1, 1)$ and so in the kernel of β . It therefore comes from a unique holomorphic line bundle $\mathcal{L}_t \in H^1(\mathcal{X}_t, \mathcal{O}^*)$. Moreover, we can always choose a representative $(U, 0)$ of the germ $(\text{Def}(X), 0)$ such that $U \cap \Delta^f$ is connected and simply-connected, so there exists a line bundle $\mathcal{L}$ over $\mathcal{X}|_{\Delta^f}$, restricting to $\mathcal{L}_t$ on each fibre $\mathcal{X}_t$.

By Theorem 3, $h^i(\mathcal{X}_t, \mathcal{L}_t)$ is an upper semi-continuous function of $t \in \Delta^f$. So for t in a neighbourhood of zero

$$h^i(\mathcal{X}_t, \mathcal{L}_t) \leq h^i(\mathcal{X}_0, \mathcal{L}_0).$$

But the right hand side is $h^i(X, L)$, which by Lemma 2 is $n + 1$ for $i = 0$ and zero otherwise. Therefore

$$h^i(\mathcal{X}_t, \mathcal{L}_t) = 0$$

for $i > 0$. On the other hand, the Euler characteristic

$$\chi(\mathcal{X}_t, \mathcal{L}_t) := \sum_{i=0}^{2n} (-1)^i h^i(\mathcal{X}_t, \mathcal{L}_t)$$

is constant (it is given by the Hirzebruch-Riemann-Roch formula) and equal to $\chi(X, L) = n + 1$. Therefore

$$h^0(\mathcal{X}_t, \mathcal{L}_t) = n + 1$$

for all t in a neighbourhood of zero. The linear system of $\mathcal{L}_t$ therefore gives a map

$$\phi_t : \mathcal{X}_t \rightarrow \mathbb{P}(H^0(\mathcal{X}_t, \mathcal{L}_t)^*) \cong \mathbb{P}^n.$$

A priori ϕ_t is only a rational map, so we must show that the linear system of $\mathcal{L}_t$ has no base-points. This follows since the specialization

$$\phi_L = f : X \rightarrow \mathbb{P}(H^0(X, L)^*) \cong \mathbb{P}^n$$

is a morphism and thus has no base-points. More specifically, we can extend a basis $\{s_0, \dots, s_n\}$ of sections of L to a set of $n + 1$ sections $\{S_0, \dots, S_n\}$ of $\mathcal{L}$ over $\mathcal{X}|_{\Delta^f} \rightarrow \Delta^f$. The locus

$$C := \{t \in \Delta^f \mid S_0|_{\mathcal{X}_t}, \dots, S_n|_{\mathcal{X}_t} \text{ simultaneously vanish at some point of } \mathcal{X}_t\}$$

is a closed subset of Δ^f which does not contain 0. So for $t \in \Delta^f$ in a neighbourhood of zero, the sections $S_0|_{\mathcal{X}_t}, \dots, S_n|_{\mathcal{X}_t}$ generate the fibres of $\mathcal{L}_t$.

We have proved that

$$\phi_t : \mathcal{X}_t \rightarrow \mathbb{P}^n$$

is a morphism, and it is therefore a Lagrangian fibration by the results of Matsushita (and Huybrechts) cited in the introduction. $\square$

4 Fibrations with sections

We will use the following result (note that in Voisin's paper X is the Lagrangian submanifold of Y).

Theorem 5 (Voisin [16]) *Let X be a holomorphic symplectic manifold and $g : Y \hookrightarrow X$ a Lagrangian submanifold. Let g_t^* be the composition*

$$g^* \circ (\psi_t \otimes \mathbb{C}) : H^2(\mathcal{X}_t, \mathbb{C}) \rightarrow H^2(Y, \mathbb{C})$$

where ψ_t is a marking of $\mathcal{X}_t$. Then the inclusion $Y \hookrightarrow X$ deforms to a Lagrangian submanifold $\mathcal{Y}_t \rightarrow \mathcal{X}_t$ if and only if $g_t^*(\sigma_t) = 0$.

We can rephrase this as follows. Let $L_Y \subset H^2(X, \mathbb{C})$ be the orthogonal complement of $\ker g^*$ with respect to the Beauville-Bogomolov quadratic form q_X . Since $H^{2,0}(X) \subset \ker g^*$ and L_Y can be defined over $\mathbb{Q}$, L_Y must be of type $(1, 1)$. Then $Y \hookrightarrow X$ deforms to $\mathcal{Y}_t \rightarrow \mathcal{X}_t$ if and only if L_Y is preserved under the deformation, i.e. if and only if

$$t \in \Delta := \{t \in \text{Def}(X) \mid \psi_t^{-1}(L_Y) \in H^2(\mathcal{X}_t, \mathbb{C}) \text{ is of type } (1, 1)\}.$$

We wish to look at deformations of a Lagrangian fibration $f : X \rightarrow B$ which admits a section s . Note that the existence of the section implies that $B \cong \mathbb{P}^n$ [3]. However, we will keep the projectivity of X as a hypothesis. Let Y denote the image of s , and $g : Y \hookrightarrow X$ the inclusion. Since Y is isomorphic to $B \cong \mathbb{P}^n$, it must be holomorphic Lagrangian, as

$$\sigma|_Y \in H^0(Y, \Omega^2) \cong H^{2,0}(\mathbb{P}^n) = 0.$$

Theorem 6 *Let X be an irreducible projective holomorphic symplectic manifold which admits a Lagrangian fibration $f : X \rightarrow B$ and a global section s . There is a codimension two submanifold $\Delta^f \subset \text{Def}(X)$ (containing zero) which parametrizes holomorphic symplectic manifolds X' which admit Lagrangian fibrations and global sections.*

Proof Firstly, the existence of the section implies $B \cong \mathbb{P}^n$, and so by Theorem 4 there exists a codimension one family $\Delta^f \subset \text{Def}(X)$ over which the fibration deforms.

By Theorem 5 we also have

$$\Delta^s := \{t \in \text{Def}(X) \mid \psi_t^{-1}(L_Y) \in H^2(\mathcal{X}_t, \mathbb{C}) \text{ is of type } (1, 1)\}$$

over which the Lagrangian submanifold $Y \hookrightarrow X$ deforms. Now Δ^s maps isomorphically to a neighbourhood of $[\sigma]$ in

$$Q_X^s := \{[\alpha] \in Q_X \mid q_X(\alpha, L_Y) = 0\} = \{[\alpha] \in Q_X \mid \alpha \in \ker g^*\}.$$

Since $Y \cong \mathbb{P}^n$, $H^2(Y, \mathbb{C})$ and L_Y are one-dimensional, so Q_X^s and Δ^s are codimension one in Q_X and $\text{Def}(X)$ respectively.

Let $\Delta^{fs} := \Delta^f \cap \Delta^s$. It maps isomorphically to a neighbourhood of $[\sigma]$ in

$$Q_X^{fs} := Q_X^f \cap Q_X^s = \{[\alpha] \in Q_X \mid q_X(\alpha, c_1) = 0 \text{ and } \alpha \in \ker g^*\}.$$

Observe that $L := f^*\mathcal{O}(1)$ must satisfy

$$c_1(L)^{n+1} = 0 \in H^{2n+2}(X, \mathbb{Z}),$$

which implies $q_X(c_1) = 0$ (see [7]) and thus $[c_1] \in Q_X^f$. On the other hand

$$g^*(c_1) = c_1(\mathcal{O}(1)) \neq 0,$$

so $c_1 \notin \ker g^*$ and $[c_1] \notin Q_X^s$. This suffices to show that Q_X^f and Q_X^s intersect transversely, and thus Q_X^{fs} and Δ^{fs} are codimension two in Q_X and $\text{Def}(X)$ respectively.

Finally observe that if $t \in \Delta^{fs}$ then $\mathcal{X}_t$ is a Lagrangian fibration over $\mathbb{P}^n$ and it contains a deformation $\mathcal{Y}_t$ of Y . By specialization, $\mathcal{Y}_t$ is a section of the fibration. More precisely, the set of values t for which $\mathcal{Y}_t$ maps isomorphically to $\mathbb{P}^n$ is open in Δ^{fs} and contains zero. This completes the proof. $\square$

Example Let S be a K3 surface which contains a smooth genus g curve C , $g \geq 2$, such that $\text{Pic}S$ is generated by $[C]$. Let $\mathcal{C} \rightarrow |C| \cong \mathbb{P}^g$ be the family of curves linearly equivalent to C . Then the compactified Jacobian

$$J^0 := \overline{\text{Jac}}^0(\mathcal{C}/\mathbb{P}^g)$$

is a deformation of the Hilbert scheme of g points on S [2]. Moreover, J^0 is a Lagrangian fibration over $\mathbb{P}^g$ which admits a section. Observe that we can deform S in a 19-dimensional family, while keeping the genus g curve: these produce deformations of J^0 keeping the Lagrangian fibration and section. On the other hand, the space of deformations of the Hilbert scheme of points on S is 21-dimensional. This is in agreement with Theorem 6.

We find similar agreement for Lagrangian fibrations which are deformations of the generalized Kummer varieties (see Debarre [4], or the author's paper [13]).

Remark Let $f : X \rightarrow B$ be a Lagrangian fibration. If the fibres of X are all reduced and irreducible then there exists the (Altman-Kleiman) compactified relative Picard scheme

$$P := \overline{\text{Pic}}^0(X/B).$$

If the fibres of P are reduced and irreducible we also have

$$X^0 := \overline{\text{Pic}}^0(P/B).$$

Both P and X^0 admit global sections. Let U and U^0 denote the open subsets of X and X^0 , respectively, given by removing all singular fibres; then U is a torsor over U^0 . In fact in cases where the singular fibres of X are not too complicated, X itself is a torsor over X^0 ; moreover P and X^0 are holomorphic symplectic manifolds (see [15]). In [14] it was shown that torsors X over X^0 are parametrized by the one-dimensional space $H^2(P, \mathcal{O}^*)$ of gerbes on P , with X^0 being the unique fibration which admits a global section. This agrees with Theorem 4 and Theorem 6, which together imply that fibrations which admit sections must be codimension one inside the family of all fibrations.

References

- [1] A. Beauville, *Variétés Kähleriennes dont le 1^{ère} classe de Chern est nulle*, Jour. Diff. Geom. **18** (1983), 755–782.
- [2] A. Beauville, *Counting rational curves on K3 surfaces*, Duke Math. J. **97** (1999), no. 1, 99–108.
- [3] K. Cho, Y. Miyaoka, and N. Shepherd-Barron, *Characterizations of projective space and applications to complex symplectic manifolds*, in Higher Dimensional Birational Geometry, Advanced Studies in Pure Mathematics **35**, 2002, 1–88.
- [4] O. Debarre, *On the Euler characteristic of generalized Kummer varieties*, Amer. J. Math. **121** (1999), no. 3, 577–586.
- [5] M. Eastwood and J. Sawon, *The Borel-Weil theorem for complex projective space*, in Invitations to Geometry and Topology, Oxford University Press (2002), 126–145.
- [6] H. Grauert and R. Remmert, *Coherent analytic sheaves*, Grundlehren der Mathematischen Wissenschaften **265**, Springer-Verlag, Berlin, 1984.
- [7] M. Gross, D. Huybrechts, and D. Joyce, *Calabi-Yau manifolds and related geometries*, Springer Universitext, 2002.
- [8] D. Markushevich, *Lagrangian families of Jacobians of genus 2 curves*, J. Math. Sci. **82** (1996), no. 1, 3268–3284.
- [9] D. Matsushita, *On fibre space structures of a projective irreducible symplectic manifold*, Topology **38** (1999), No. 1, 79–83. Addendum, Topology **40** (2001), no. 2, 431–432.
- [10] D. Matsushita, *Equidimensionality of Lagrangian fibrations on holomorphic symplectic manifolds*, Math. Res. Lett. **7** (2000), no. 4, 389–391.
- [11] D. Matsushita, *Higher direct images of dualizing sheaves of Lagrangian fibrations*, Amer. J. Math. **127** (2005), no. 2, 243–259.
- [12] A. Rapagnetta, *Topological invariants of O’Grady’s six dimensional irreducible symplectic variety*, preprint **math.AG/0406026**.
- [13] J. Sawon, *Abelian fibred holomorphic symplectic manifolds*, Turkish Jour. Math. **27** (2003), no. 1, 197–230.
- [14] J. Sawon, *Derived equivalence of holomorphic symplectic manifolds*, in Algebraic Structures and Moduli Spaces, CRM Proc. Lecture Notes **38**, AMS, 2004, 193–211.
- [15] J. Sawon, *Twisted Fourier-Mukai transforms for holomorphic symplectic four-folds*, preprint.

- [16] C. Voisin, *Sur la stabilité des sous-variétés lagrangiennes des variétés symplectiques holomorphes*, Complex projective geometry, 294–303, LMS Lecture Note Ser. **179**, Cambridge Univ. Press, 1992.

Department of Mathematics
SUNY at Stony Brook
Stony Brook NY 11794-3651
USA

sawon@math.sunysb.edu
www.math.sunysb.edu/~sawon